

FLURALANER INJECTABLE – Name brand: Bravecto Quantum

GENERAL DESCRIPTION:

Fluralaner Injectable is a prescribed drug that has activity against adult fleas and certain tick species. Fleas live on your pet causing discomfort and skin irritation while ticks can transmit serious diseases and may pose a health risk to other household members. Fluralaner is available as an extended-release injectable solution for dogs administered by a licensed veterinarian that may provide 8-12 months of protection based on local tick species.

WHAT IS THIS DRUG?

- Fluralaner is a long-acting ectoparasiticide (insecticide and acaricide) that belongs to the isoxazoline group
- Fluralaner Injectable has sustained activity against adult fleas for 12 months in dogs
- Fluralaner Injectable has sustained activity against the Black-legged tick, American dog tick, Brown Dog tick for 12 months in dogs
- Fluralaner Injectable has sustained activity against the Lone Star tick for 8 months in dogs
- Given by injection under the skin (SQ) every 12 months (every 8 months in areas endemic with the Lone Star tick)

REASONS FOR PRESCRIBING:

- Used for the treatment and prevention of adult fleas
- Used for the treatment and control of certain tick species

ANIMAL PRECAUTIONS:

- This medication is labeled for use in DOGS and should be administered only when and as prescribed by a licensed veterinarian
- Use of this medication in animals with any of the following conditions should be limited to those who are under the supervision of a licensed veterinarian:
 - Previous history of seizures or underlying neurological disorders
- Safety has not been established in:

- Puppies less than 6 months of age and weighing less than 2.2 lb (1 kg)
 - Breeding, pregnant or nursing animals
- Animals with a known hypersensitivity or allergy to this or related products should not be given this medication

DIRECTIONS:

This injectable medication is given in the safety of a veterinary hospital with adequate patient monitoring.

This medication should take effect within 3-5 days. You should see a notable reduction in parasites.

If you have a flea infestation in your home or your other pets are not on regular flea prevention, you may see reinfestation of your pet with fleas within the first 3 months. Home extermination of fleas and treatment of all pets in the household for fleas is essential for any flea product to work effectively.

WHAT IF A DOSE IS MISSED?

If an injection is missed, schedule an additional appointment at your earliest convenience.

WHAT TO TELL/ASK YOUR VETERINARIAN BEFORE GIVING THIS MEDICATION?

Talk to your veterinarian about:

- The life cycle of the flea
- How to safely remove an attached tick
- What are the risks and benefits of using this product

Tell your veterinarian:

- If your pet has experienced side effects on other drugs/products
- If your pet has experienced digestive upset now or ever
- If your pet has experienced liver or kidney disease now or ever
- If your pet has experienced any other medical problems or allergies now or ever
- All medicines and supplements that you are giving your pet or plan to give your pet, including those you can get without a

prescription. Your veterinarian may want to check that all of your pet's medications can be given together.

- If your pet is pregnant, nursing or if you plan to breed your pet

POTENTIAL SIDE EFFECTS ARE VERY RARE, BUT CAN INCLUDE:

- This medication can affect the gastrointestinal system causing a decrease in appetite, vomiting and diarrhea. It can affect the neurological system causing lethargy, muscle tremors, incoordination (ataxia) and seizures
- Injection site reaction may include swelling, pain, redness, bleeding or abscess formation
- This medication can cause pruritus, dermatitis, immune mediated disorders (anaphylaxis and immune mediated hemolytic anemia)
- This medication has the potential to cause an immediate allergic or anaphylactic reaction. Affected animals may have vomiting, diarrhea, swelling, hives, changes in respiration and display signs of agitation. This can be a severe and life-threatening; immediate veterinary care is often required.
- Adverse events have been reported in breeding females including birth defects (limb deformities and cleft palate), stillbirths and abortions
- If you notice any of these symptoms or if your pet's condition worsens, contact your veterinarian

CAN THIS DRUG BE GIVEN WITH OTHER MEDICATIONS?

- Ensure that your veterinarian knows all the medicines and supplements that you are giving your pet or plan to give your pet, including those you can get without a prescription.
- If your pet experiences any unusual reactions when taking multiple medications, contact your veterinarian.

WHAT SHOULD I DO IF I KNOW OF OR SUSPECT THERE HAS BEEN AN OVERDOSE?

Contact your veterinarian and/or a poison control center immediately if your pet receives more than the prescribed amount.

WHAT ELSE SHOULD I KNOW?

Notify your veterinarian if your animal's condition does not improve or worsens despite this treatment.

This product is not effective against internal parasites.

As with all prescribed medication, Fluralaner Injectable should only be given to the pet for which it was prescribed. It should be given only for the condition for which it was prescribed, at the prescribed dose, as directed by a veterinarian.

When dealing with a flea infestation, it is important to treat all animals within the household and the environment. Speak to your veterinarian about approved and safe products for all household pets. If fleas are already developing in your house, ask your veterinarian to recommend a premises control product as well.